arXiv:1010.1246v1 [nucl-ex] 6 Oct 2010

Cold Nuclear Matter Effects on J/ψ Yields as a Function of Rapidity and Nuclear Geometry in d+A Collisions at $\sqrt{s_{NN}}$=200 GeV

A. Adare,[12] S. Afanasiev,[27] C. Aidala,[40] N.N. Ajitanand,[57] Y. Akiba,[51,52] H. Al-Bataineh,[46] J. Alexander,[57]
A. Angerami,[13] K. Aoki,[33,51] N. Apadula,[58] L. Aphecetche,[59] Y. Aramaki,[11] J. Asai,[51] E.T. Atomssa,[34]
R. Averbeck,[58] T.C. Awes,[47] B. Azmoun,[6] V. Babintsev,[23] M. Bai,[5] G. Baksay,[19] L. Baksay,[19] A. Baldisseri,[15]
K.N. Barish,[7] P.D. Barnes,[36] B. Bassalleck,[45] A.T. Basye,[1] S. Bathe,[7,52] S. Batsouli,[47] V. Baublis,[50] C. Baumann,[41]
A. Bazilevsky,[6] S. Belikov,[6,*] R. Belmont,[63] R. Bennett,[58] A. Berdnikov,[54] Y. Berdnikov,[54] J.H. Bhom,[66]
A.A. Bickley,[12] D.S. Blau,[32] J.G. Boissevain,[36] J.S. Bok,[66] H. Borel,[15] K. Boyle,[58] M.L. Brooks,[36] H. Buesching,[6]
V. Bumazhnov,[23] G. Bunce,[6,52] S. Butsyk,[36] C.M. Camacho,[36] S. Campbell,[58] A. Caringi,[42] B.S. Chang,[66]
W.C. Chang,[2] J.-L. Charvet,[15] C.-H. Chen,[58] S. Chernichenko,[23] C.Y. Chi,[13] M. Chiu,[6,24] I.J. Choi,[66] J.B. Choi,[9]
R.K. Choudhury,[4] P. Christiansen,[38] T. Chujo,[62] P. Chung,[57] A. Churyn,[23] O. Chvala,[7] V. Cianciolo,[47]
Z. Citron,[58] B.A. Cole,[13] Z. Conesa del Valle,[34] M. Connors,[58] P. Constantin,[36] M. Csanád,[17] T. Csörgő,[30]
T. Dahms,[58] S. Dairaku,[33,51] I. Danchev,[63] K. Das,[20] A. Datta,[40] G. David,[6] M.K. Dayananda,[21] A. Denisov,[23]
D. d'Enterria,[34] A. Deshpande,[52,58] E.J. Desmond,[6] K.V. Dharmawardane,[46] O. Dietzsch,[55] A. Dion,[26,58]
M. Donadelli,[55] L. D Orazio,[39] O. Drapier,[34] A. Drees,[58] K.A. Drees,[5] A.K. Dubey,[65] J.M. Durham,[58]
A. Durum,[23] D. Dutta,[4] V. Dzhordzhadze,[7] S. Edwards,[20] Y.V. Efremenko,[47] F. Ellinghaus,[12] T. Engelmore,[13]
A. Enokizono,[35,47] H. En'yo,[51,52] S. Esumi,[62] K.O. Eyser,[7] B. Fadem,[42] D.E. Fields,[45,52] M. Finger, Jr.,[8]
M. Finger,[8] F. Fleuret,[34] S.L. Fokin,[32] Z. Fraenkel,[65,*] J.E. Frantz,[58] A. Franz,[6] A.D. Frawley,[20] K. Fujiwara,[51]
Y. Fukao,[33,51] T. Fusayasu,[44] I. Garishvili,[60] A. Glenn,[12,35] H. Gong,[58] M. Gonin,[34] J. Gosset,[15] Y. Goto,[51,52]
R. Granier de Cassagnac,[34] N. Grau,[13] S.V. Greene,[63] G. Grim,[36] M. Grosse Perdekamp,[24,52] T. Gunji,[11]
H.-Å. Gustafsson,[38,*] A. Hadj Henni,[59] J.S. Haggerty,[6] K.I. Hahn,[18] H. Hamagaki,[11] J. Hamblen,[60] J. Hanks,[13]
R. Han,[49] E.P. Hartouni,[35] K. Haruna,[22] E. Haslum,[38] R. Hayano,[11] M. Heffner,[35] T.K. Hemmick,[58] T. Hester,[7]
X. He,[21] J.C. Hill,[26] M. Hohlmann,[19] W. Holzmann,[13,57] K. Homma,[22] B. Hong,[31] T. Horaguchi,[11,22,51,61]
D. Hornback,[60] S. Huang,[63] T. Ichihara,[51,52] R. Ichimiya,[51] H. Iinuma,[33,51] Y. Ikeda,[62] K. Imai,[33,51] J. Imrek,[16]
M. Inaba,[62] D. Isenhower,[1] M. Ishihara,[51] T. Isobe,[11] M. Issah,[57,63] A. Isupov,[27] D. Ivanischev,[50] Y. Iwanaga,[22]
B.V. Jacak,[58,†] J. Jia,[6,13,57] X. Jiang,[36] J. Jin,[13] B.M. Johnson,[6] T. Jones,[1] K.S. Joo,[43] D. Jouan,[48] D.S. Jumper,[1]
F. Kajihara,[11] S. Kametani,[51] N. Kamihara,[52] J. Kamin,[58] J.H. Kang,[66] J. Kapustinsky,[36] K. Karatsu,[33]
M. Kasai,[53,51] D. Kawall,[40,52] M. Kawashima,[53,51] A.V. Kazantsev,[32] T. Kempel,[26] A. Khanzadeev,[50]
K.M. Kijima,[22] J. Kikuchi,[64] A. Kim,[18] B.I. Kim,[31] D.H. Kim,[43] D.J. Kim,[28,66] E.J. Kim,[9] E. Kim,[56] S.H. Kim,[66]
Y.-J. Kim,[24] E. Kinney,[12] K. Kiriluk,[12] Á. Kiss,[17] E. Kistenev,[6] J. Klay,[35] C. Klein-Boesing,[41] L. Kochenda,[50]
B. Komkov,[50] M. Konno,[62] J. Koster,[24] A. Kozlov,[65] A. Král,[14] A. Kravitz,[13] G.J. Kunde,[36] K. Kurita,[53,51]
M. Kurosawa,[51] M.J. Kweon,[31] Y. Kwon,[60,66] G.S. Kyle,[46] R. Lacey,[57] Y.S. Lai,[13] J.G. Lajoie,[26] D. Layton,[24]
A. Lebedev,[26] D.M. Lee,[36] J. Lee,[18] K.B. Lee,[31] K.S. Lee,[31] T. Lee,[56] M.J. Leitch,[36] M.A.L. Leite,[55] B. Lenzi,[55]
P. Lichtenwalner,[42] P. Liebing,[52] L.A. Linden Levy,[12] T. Liška,[14] A. Litvinenko,[27] H. Liu,[36,46] M.X. Liu,[36]
X. Li,[10] B. Love,[63] D. Lynch,[6] C.F. Maguire,[63] Y.I. Makdisi,[5] A. Malakhov,[27] M.D. Malik,[45] V.I. Manko,[32]
E. Mannel,[13] Y. Mao,[49,51] L. Mašek,[8,25] H. Masui,[62] F. Matathias,[13] M. McCumber,[58] P.L. McGaughey,[36]
D. McGlinchey,[20] N. Means,[58] B. Meredith,[24] Y. Miake,[62] T. Mibe,[29] A.C. Mignerey,[39] P. Mikeš,[25] K. Miki,[62]
A. Milov,[6] M. Mishra,[3] J.T. Mitchell,[6] A.K. Mohanty,[4] H.J. Moon,[43] Y. Morino,[11] A. Morreale,[7] D.P. Morrison,[6]
T.V. Moukhanova,[32] D. Mukhopadhyay,[63] T. Murakami,[33] J. Murata,[53,51] S. Nagamiya,[29] J.L. Nagle,[12]
M. Naglis,[65] M.I. Nagy,[17,30] I. Nakagawa,[51,52] Y. Nakamiya,[22] K.R. Nakamura,[33] T. Nakamura,[22,51]
K. Nakano,[51,61] S. Nam,[18] J. Newby,[35] M. Nguyen,[58] M. Nihashi,[22] T. Niita,[62] R. Nouicer,[6] A.S. Nyanin,[32]
C. Oakley,[21] E. O'Brien,[6] S.X. Oda,[11] C.A. Ogilvie,[26] K. Okada,[52] M. Oka,[62] Y. Onuki,[51] A. Oskarsson,[38]
M. Ouchida,[22] K. Ozawa,[11] R. Pak,[6] A.P.T. Palounek,[36] V. Pantuev,[58] V. Papavassiliou,[46] I.H. Park,[18] J. Park,[56]
S.K. Park,[31] W.J. Park,[31] S.F. Pate,[46] H. Pei,[26] J.-C. Peng,[24] H. Pereira,[15] V. Peresedov,[27] D.Yu. Peressounko,[32]
R. Petti,[58] C. Pinkenburg,[6] R.P. Pisani,[6] M. Proissl,[58] M.L. Purschke,[6] A.K. Purwar,[36] H. Qu,[21] J. Rak,[28,45]
A. Rakotozafindrabe,[34] I. Ravinovich,[65] K.F. Read,[47,60] S. Rembeczki,[19] K. Reygers,[41] V. Riabov,[50] Y. Riabov,[50]
E. Richardson,[39] D. Roach,[63] G. Roche,[37] S.D. Rolnick,[7] M. Rosati,[26] C.A. Rosen,[12] S.S.E. Rosendahl,[38]
P. Rosnet,[37] P. Rukoyatkin,[27] P. Ružička,[25] V.L. Rykov,[51] B. Sahlmueller,[41] N. Saito,[29,33,51,52] T. Sakaguchi,[6]
S. Sakai,[62] K. Sakashita,[51,61] V. Samsonov,[50] S. Sano,[11,64] T. Sato,[62] S. Sawada,[29] K. Sedgwick,[7] J. Seele,[12]
R. Seidl,[24,52] A.Yu. Semenov,[26] V. Semenov,[23] R. Seto,[7] D. Sharma,[65] I. Shein,[23] T.-A. Shibata,[51,61] K. Shigaki,[22]

M. Shimomura,[62] K. Shoji,[33, 51] P. Shukla,[4] A. Sickles,[6] C.L. Silva,[26, 55] D. Silvermyr,[47] C. Silvestre,[15] K.S. Sim,[31] B.K. Singh,[3] C.P. Singh,[3] V. Singh,[3] M. Slunečka,[8] A. Soldatov,[23] R.A. Soltz,[35] W.E. Sondheim,[36] S.P. Sorensen,[60] I.V. Sourikova,[6] F. Staley,[15] P.W. Stankus,[47] E. Stenlund,[38] M. Stepanov,[46] A. Ster,[30] S.P. Stoll,[6] T. Sugitate,[22] C. Suire,[48] A. Sukhanov,[6] J. Sziklai,[30] E.M. Takagui,[55] A. Taketani,[51, 52] R. Tanabe,[62] Y. Tanaka,[44] S. Taneja,[58] K. Tanida,[33, 51, 52, 56] M.J. Tannenbaum,[6] S. Tarafdar,[3] A. Taranenko,[57] P. Tarján,[16] H. Themann,[58] D. Thomas,[1] T.L. Thomas,[45] M. Togawa,[33, 51, 52] A. Toia,[58] L. Tomášek,[25] Y. Tomita,[62] H. Torii,[22, 51] R.S. Towell,[1] V-N. Tram,[34] I. Tserruya,[65] Y. Tsuchimoto,[22] C. Vale,[6, 26] H. Valle,[63] H.W. van Hecke,[36] E. Vazquez-Zambrano,[13] A. Veicht,[24] J. Velkovska,[63] R. Vértesi,[16, 30] A.A. Vinogradov,[32] M. Virius,[14] A. Vossen,[24] V. Vrba,[25] E. Vznuzdaev,[50] X.R. Wang,[46] D. Watanabe,[22] K. Watanabe,[62] Y. Watanabe,[51, 52] F. Wei,[26] R. Wei,[57] J. Wessels,[41] S.N. White,[6] D. Winter,[13] C.L. Woody,[6] R.M. Wright,[1] M. Wysocki,[12] W. Xie,[52] Y.L. Yamaguchi,[11, 64] K. Yamaura,[22] R. Yang,[24] A. Yanovich,[23] J. Ying,[21] S. Yokkaichi,[51, 52] G.R. Young,[47] I. Younus,[45] Z. You,[49] I.E. Yushmanov,[32] W.A. Zajc,[13] O. Zaudtke,[41] C. Zhang,[47] S. Zhou,[10] and L. Zolin[27]

(PHENIX Collaboration)

[1]*Abilene Christian University, Abilene, Texas 79699, USA*
[2]*Institute of Physics, Academia Sinica, Taipei 11529, Taiwan*
[3]*Department of Physics, Banaras Hindu University, Varanasi 221005, India*
[4]*Bhabha Atomic Research Centre, Bombay 400 085, India*
[5]*Collider-Accelerator Department, Brookhaven National Laboratory, Upton, New York 11973-5000, USA*
[6]*Physics Department, Brookhaven National Laboratory, Upton, New York 11973-5000, USA*
[7]*University of California - Riverside, Riverside, California 92521, USA*
[8]*Charles University, Ovocný trh 5, Praha 1, 116 36, Prague, Czech Republic*
[9]*Chonbuk National University, Jeonju, 561-756, Korea*
[10]*China Institute of Atomic Energy (CIAE), Beijing, People's Republic of China*
[11]*Center for Nuclear Study, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan*
[12]*University of Colorado, Boulder, Colorado 80309, USA*
[13]*Columbia University, New York, New York 10027 and Nevis Laboratories, Irvington, New York 10533, USA*
[14]*Czech Technical University, Zikova 4, 166 36 Prague 6, Czech Republic*
[15]*Dapnia, CEA Saclay, F-91191, Gif-sur-Yvette, France*
[16]*Debrecen University, H-4010 Debrecen, Egyetem tér 1, Hungary*
[17]*ELTE, Eötvös Loránd University, H - 1117 Budapest, Pázmány P. s. 1/A, Hungary*
[18]*Ewha Womans University, Seoul 120-750, Korea*
[19]*Florida Institute of Technology, Melbourne, Florida 32901, USA*
[20]*Florida State University, Tallahassee, Florida 32306, USA*
[21]*Georgia State University, Atlanta, Georgia 30303, USA*
[22]*Hiroshima University, Kagamiyama, Higashi-Hiroshima 739-8526, Japan*
[23]*IHEP Protvino, State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, 142281, Russia*
[24]*University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA*
[25]*Institute of Physics, Academy of Sciences of the Czech Republic, Na Slovance 2, 182 21 Prague 8, Czech Republic*
[26]*Iowa State University, Ames, Iowa 50011, USA*
[27]*Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia*
[28]*Helsinki Institute of Physics and University of Jyväskylä, P.O.Box 35, FI-40014 Jyväskylä, Finland*
[29]*KEK, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan*
[30]*KFKI Research Institute for Particle and Nuclear Physics of the Hungarian Academy of Sciences (MTA KFKI RMKI), H-1525 Budapest 114, POBox 49, Budapest, Hungary*
[31]*Korea University, Seoul, 136-701, Korea*
[32]*Russian Research Center "Kurchatov Institute", Moscow, Russia*
[33]*Kyoto University, Kyoto 606-8502, Japan*
[34]*Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS-IN2P3, Route de Saclay, F-91128, Palaiseau, France*
[35]*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*
[36]*Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*
[37]*LPC, Université Blaise Pascal, CNRS-IN2P3, Clermont-Fd, 63177 Aubiere Cedex, France*
[38]*Department of Physics, Lund University, Box 118, SE-221 00 Lund, Sweden*
[39]*University of Maryland, College Park, Maryland 20742, USA*
[40]*Department of Physics, University of Massachusetts, Amherst, Massachusetts 01003-9337, USA*
[41]*Institut fur Kernphysik, University of Muenster, D-48149 Muenster, Germany*
[42]*Muhlenberg College, Allentown, Pennsylvania 18104-5586, USA*
[43]*Myongji University, Yongin, Kyonggido 449-728, Korea*
[44]*Nagasaki Institute of Applied Science, Nagasaki-shi, Nagasaki 851-0193, Japan*
[45]*University of New Mexico, Albuquerque, New Mexico 87131, USA*
[46]*New Mexico State University, Las Cruces, New Mexico 88003, USA*

[47]*Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA*
[48]*IPN-Orsay, Universite Paris Sud, CNRS-IN2P3, BP1, F-91406, Orsay, France*
[49]*Peking University, Beijing, People's Republic of China*
[50]*PNPI, Petersburg Nuclear Physics Institute, Gatchina, Leningrad region, 188300, Russia*
[51]*RIKEN Nishina Center for Accelerator-Based Science, Wako, Saitama 351-0198, Japan*
[52]*RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, New York 11973-5000, USA*
[53]*Physics Department, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima, Tokyo 171-8501, Japan*
[54]*Saint Petersburg State Polytechnic University, St. Petersburg, Russia*
[55]*Universidade de São Paulo, Instituto de Física, Caixa Postal 66318, São Paulo CEP05315-970, Brazil*
[56]*Seoul National University, Seoul, Korea*
[57]*Chemistry Department, Stony Brook University, SUNY, Stony Brook, New York 11794-3400, USA*
[58]*Department of Physics and Astronomy, Stony Brook University, SUNY, Stony Brook, New York 11794-3400, USA*
[59]*SUBATECH (Ecole des Mines de Nantes, CNRS-IN2P3, Université de Nantes) BP 20722 - 44307, Nantes, France*
[60]*University of Tennessee, Knoxville, Tennessee 37996, USA*
[61]*Department of Physics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan*
[62]*Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
[63]*Vanderbilt University, Nashville, Tennessee 37235, USA*
[64]*Waseda University, Advanced Research Institute for Science and*
Engineering, 17 Kikui-cho, Shinjuku-ku, Tokyo 162-0044, Japan
[65]*Weizmann Institute, Rehovot 76100, Israel*
[66]*Yonsei University, IPAP, Seoul 120-749, Korea*

(Dated: October 7, 2010)

We present measurements of J/ψ yields in d+Au collisions at $\sqrt{s_{_{NN}}}$=200 GeV recorded by the PHENIX experiment and compare with yields in $p+p$ collisions at the same energy per nucleon-nucleon collision. The measurements cover a large kinematic range in J/ψ rapidity ($-2.2 < y < 2.4$) with high statistical precision and are compared with two theoretical models: one with nuclear shadowing combined with final state breakup and one with coherent gluon saturation effects. In order to remove model dependent systematic uncertainties we also compare the data to a simple geometric model. We find that calculations where the nuclear modification is linear or exponential in the density weighted longitudinal thickness are difficult to reconcile with the forward rapidity data.

PACS numbers: 25.75.Dw

The measured yields of quarkonia states in p+A (or d+A) collisions provide information about the time scale and dynamics for the creation of a $c\overline{c}$ pair and its evolution to a color-singlet quarkonium state. The propagation time of the $c\overline{c}$ pair through the nucleus is set by the incident energy of the projectile and target and by the relative longitudinal momentum of the $c\overline{c}$ pair. Fixed target p+A experiments at Fermilab [1] reveal a substantial suppression for forward rapidity J/ψ and ψ' at a similar level, leading to the conclusion that the suppression must occur at the prehadronic stage. An analysis [2] of results for $\sqrt{s_{_{NN}}}$ = 17–42 GeV indicates that in addition to modified initial production due to nuclear-modified parton distribution functions (nPDFs), a break up cross section (σ_{br}) for the $c\overline{c}$ precursor state to the J/ψ is important, and that σ_{br} decreases as the relative center-of-mass energy between the $c\overline{c}$ and the nucleon increases. Extending these results to collider energies at RHIC is important. The dominant production mechanism for charm (at RHIC) is via gluon-gluon interactions, and thus the yields at forward rapidity, the deuteron-going direction, are sensitive to low-x in the gold nucleus where gluon shadowing [3, 4] and/or gluon saturation effects [5] become important, providing a crucial test of these effects.

There is also significant interest in determining the color screening length in the quark-gluon plasma for temperatures $T > 170$ MeV, as achieved in relativistic heavy ion collisions [6]. One proposal for determining this is the measurement of several quarkonia states where the different binding energies (and thus radii) can bracket the screening length of interest [7, 8]. However, this suppression of quarkonia must be separated from the aforementioned cold nuclear matter effects. Thus precise measurement of quarkonia suppression in d+Au is needed.

The PHENIX experiment has previously published J/ψ results in d+Au collisions at $\sqrt{s_{_{NN}}}$ = 200 GeV [9] from data taken in 2003. In this paper we present results from d+Au collision data taken in 2008, representing an increase in yield by a factor of 30–50 over the previous results and a reduction in the systematic uncertainties by up to a factor of two. Additionally, the $p+p$ reference data sets are updated to include higher statistics data from 2006 and 2008.

The PHENIX apparatus is described in detail in [10]. It consists of two sets of spectrometers referred to as the central arms, which measure single-particles emitted over pseudorapidity ($|\eta| < 0.35$), and the muon arms, measuring single muons over pseudorapidities ($1.2 < |\eta| < 2.4$). J/ψ particles are measured via their dielectron (dimuon) decays at mid (backward and forward) rapidities, and detailed analysis methods are given in [9, 11]. The d+Au data used for this analysis were recorded using selective Level-1 triggers in coincidence with a minimum bias interaction requirement, which requires one hit in each of two beam-beam counters (BBCs) located at positive and negative pseudorapidity ($3 < |\eta| < 3.9$). This minimum bias selection covers $88 \pm 4\%$ of the total d+Au inelastic cross section of 2260 mb [12]. This can be corrected to an unbiased sample, 100% of the total cross section, by applying a bias correction factor ($c = 0.89$) to the particle yield measured in any minimum bias event. Additional Level-1 triggers independently require (1) one hit above threshold (600 or 800 MeV) in the Electromagnetic Calorimeter with a matching hit in the Ring Imaging Čerenkov Detector identified as an electron or (2) two tracks identified as muon candidates [9]. The data sets sampled via the Level-1 triggers represent analyzed integrated luminosities for the different spectrometers of 54.0 nb^{-1} to 69.3 nb^{-1}. We use $p+p$ reference data for the midrapidity dielectrons from [13]. For the forward and backward rapidity dimuons, we report here new p+p data from 2006 and 2008 with a total integrated luminosity of 5.1 pb^{-1}.

The p_T-integrated J/ψ yield as a function of rapidity is calculated via:

$$B_{ll}\frac{dN}{dy} = \frac{cN_{J/\psi}}{N_{\rm MB}\epsilon A \Delta y} \qquad (1)$$

where B_{ll} is the branching fraction for $J/\psi \to e^+e^-$ or $\mu^+\mu^-$, $N_{J/\psi}$ is the number of J/ψ counts, c is the bias correction factor, $N_{\rm MB}$ is the number of sampled minimum bias events, Δy is the width of the rapidity bin, and ϵA represents the product of the efficiency and acceptance corrections, including the Level-1 trigger efficiency. The number of J/ψ particles is determined using the invariant mass distribution of unlike-sign lepton pairs. Approximately 38000, 8900, and 42000 J/ψ counts are measured at backward, mid, and forward rapidity, respectively. Figure 1a shows the J/ψ yields in p+p and d+Au unbiased collisions.

We quantify the cold nuclear matter effects by calculating the nuclear modification factor $R_{d\rm Au}$ as given by:

$$R_{d\rm Au}(i) = \frac{\frac{dN^{d+\rm Au}(i)}{dy}}{\langle N_{\rm coll}(i)\rangle \frac{dN^{p+p}}{dy}} \qquad (2)$$

where i is the index of the centrality bin and $\langle N_{\rm coll}(i)\rangle$ is the average number of nucleon-nucleon collisions and is determined from the total energy deposited in the BBC located at negative rapidity. For a given centrality bin $\langle N_{\rm coll}(i)\rangle$ is derived using a Glauber calculation coupled to a simulation of the BBC response (as described in [9]). The centrality bins used in this analysis are characterized as follows: central $\langle N_{\rm coll}(0\text{–}20\%)\rangle = 15.1 \pm 1.0$, $\langle N_{\rm coll}(20\text{–}40\%)\rangle = 10.3 \pm 0.7$, $\langle N_{\rm coll}(40\text{–}60\%)\rangle = 6.6 \pm 0.6$, $\langle N_{\rm coll}(60\text{–}88\%)\rangle = 3.2 \pm 0.2$ and unbiased $\langle N_{\rm coll}(0\text{–}100\%)\rangle = 7.6 \pm 0.3$. Figure 1b shows the $R_{d\rm Au}$ corresponding to unbiased collisions. Figure 2a (b) shows $R_{d\rm Au}$ corresponding to d+Au centralities of 60–88% (0–20%). Note that more central collisions correspond to cases where the nucleons in the deuteron strike closer to the middle of the gold nucleus, and thus nuclear effects are expected to be enhanced (which is seen in the data).



FIG. 1: (color online) (a) J/ψ B_{ll} dN/dy in p+p and d+Au collisions as a function of rapidity. The d+Au yields are divided by the average number of nucleon-nucleon collisions $\langle N_{\rm coll}(0\text{-}100\%)\rangle = 7.6$. (error lines) point-to-point uncorrelated uncertainties; (boxes) point-to-point correlated uncertainties; (text) global normalization scale uncertainties. (b) $R_{d\rm Au}$ nuclear modification factors for unbiased collisions.

The peripheral $R_{d\rm Au}$ favors some suppression at all rapidities, though this result is tempered by the current systematics of approximately ±15%. The central $R_{d\rm Au}$ indicates a much larger suppression for J/ψ at forward rapidity.

We also calculate the ratio $R_{\rm CP}$ as the nuclear modification between central and peripheral d+Au collision classes of events:

$$R_{\rm CP} = \frac{\frac{dN^{d+\rm Au}(0\text{–}20\%)}{dy} / \langle N_{\rm coll}(0\text{–}20\%)\rangle}{\frac{dN^{d+\rm Au}(60\text{–}88\%)}{dy} / \langle N_{\rm coll}(60\text{–}88\%)\rangle} \qquad (3)$$

Figure 2c shows the $R_{\rm CP}$ ratio for the most central category relative to the peripheral 60–88% category as a function of rapidity. The quantity $R_{\rm CP}$ has the advantage that many of the systematic uncertainties cancel in the ratio. One observes a dramatic suppression of forward rapidity yields for central d+Au events compared to peripheral events. At backward rapidity, there is little to no modification seen.



FIG. 2: (color online) Nuclear suppression factors ($R_{d\mathrm{Au}}$) for (a) peripheral, (b) central, and (c) R_{CP} as a function of rapidity.

We confront two specific calculations available in the literature with our data. The first class of calculations that are often employed include nuclear-modified PDFs and an effective σ_{br}. Here we utilize the EPS09 nuclear modified PDF set [14] and a σ_{br} = 4 mb (chosen to match the unbiased backward rapidity $R_{d\mathrm{Au}}$ data) [15]. We find a reasonable agreement within uncertainties with the unbiased $R_{d\mathrm{Au}}$ data shown in Fig. 1b. We also show as red dashed lines the differences within the EPS09 nPDFs for a single parameter change that gives the largest variation [14]. However, if one plots the same calculations for R_{CP} as shown in Fig. 2c, one sees reasonable agreement at backward and midrapidities, but a significant underprediction of the suppression for forward rapidity J/ψ in the most central events.

A second class of calculations incorporates gluon saturation effects at small-x [5, 16], and is compared with experimental data in Figs. 1-2. A modest J/ψ enhancement is predicted at midrapidity due to double-gluon exchange processes (not seen in the data) and a substantial J/ψ suppression at forward rapidity and in more central d+Au events due to saturation effects (in agreement with the data).

In order to further explore the centrality dependence of the nuclear effects we categorize each d+Au centrality class in terms of the distribution of transverse radial positions (r_T) of the nucleon-nucleon collisions relative to the center of the gold nucleus. The r_T distributions for the four centrality categories are shown in Fig. 3a. We expect that the nuclear effects are dependent on the density weighted longitudinal thickness through the gold nucleus $\left(\Lambda(r_T) \equiv \frac{1}{\rho_0}\int dz\rho(z, r_T)\right)$, where ρ_0 is the density in the middle of the nucleus. This quantity is also shown in Fig. 3a as a function of r_T.

We now posit three different functional dependencies of the nuclear modification on $\Lambda(r_T)$.

$$\text{Exponential}: M(r_T) = e^{-a\Lambda(r_T)} \tag{4}$$

$$\text{Linear}: M(r_T) = 1.0 - a\Lambda(r_T) \tag{5}$$

$$\text{Quadratic}: M(r_T) = 1.0 - a\Lambda(r_T)^2, \tag{6}$$

where a is a parameter depending on the average level of modification. The EPS09 nPDF based calculation, shown in Figs. 1 2, assumes the linear relation [15, 17] in Eq. 5 in order to make centrality-dependent predictions. In contrast, contributions from a break up of the $c\bar{c}$ via a σ_{br} follow the exponential relation in Eq. 4.

Figure 3b shows the nuclear modification R_{CP} in the most central bin versus the (unbiased) average modification $R_{d\mathrm{Au}}$. This particular set of quantities is chosen because for each of the three geometric dependencies (Eqs. 4–6), a given value of the parameter a results in a unique point on the plot and varying the parameter a results in a unique locus of points on which any suppression with that geometric dependence must lie.

The experimental data is also plotted in Fig. 3b for the same quantities. The ellipses represent a one standard deviation contour for the systematic uncertainties, which are largely uncorrelated between the unbiased $R_{d\mathrm{Au}}$ and R_{CP}. There is a substantial deviation between the exponential and linear cases and the experimental data at forward rapidity, while at mid and backward rapidities the data cannot discriminate between the cases. Thus, the forward rapidity data suggest that the dependence on $\Lambda(r_T)$ is non-linear and closer to quadratic. If the dominant physics leading to the modification is different at different rapidities, it is possible for example that the modification at backward rapidities is linear while at forward rapidities it is not. This is reinforced by the EPS09 plus σ_{br} calculation where regardless of the variation of the nPDF or σ_{br} one cannot simultaneously describe the full centrality dependence of the data as seen in Fig. 2.

Other non-linear density effects (*e.g.*, quadratic) for the geometric dependence [18] and for break up of the $c\bar{c}$ after production [19, 20] have been proposed. An alternative explanation is that initial-state parton energy



FIG. 3: (color online) (a) Normalized to unity at the maximum bin are (solid curves) transverse radial r_T distributions in the gold nucleus for four d+Au centrality selections and (dashed curve) density weighted longitudinal thickness as a function of r_T ($\Lambda(r_T)$). (b) (points) Unbiased $R_{d\mathrm{Au}}$ versus R_{CP} for the experimental data and (curves) constraint lines for three geometric dependencies of the nuclear modification.

loss results in a backward shift of the J/ψ rapidity distribution [21]. It has been observed [22] that the nuclear modification as a function of center-of-mass rapidity is similar to that observed at lower energies [1] with a steep increase in suppression at forward rapidities, as predicted for initial-state parton energy loss.

In summary, we have presented precision data on J/ψ yields in d+Au and p+p collisions at $\sqrt{s_{NN}} = 200$ GeV over a broad range in rapidity and d+Au centrality. Nuclear modification factors at forward rapidity as a function of centrality cannot be reconciled with a picture of cold nuclear matter effects (nPDFs and a σ_{br}) when an exponential or linear dependence on the nuclear thickness is employed. Effects of gluon saturation may play an important role in understanding the forward rapidity modifications, though other explanations involving initial-state parton energy loss need further investigation.

We thank the staff of the Collider-Accelerator and Physics Departments at BNL for their vital contributions. We also thank Ramona Vogt and Kirill Tuchin for useful discussions and theoretical calculations. We acknowledge support from the Office of Nuclear Physics in DOE Office of Science, NSF, and a sponsored research grant from Renaissance Technologies (USA), MEXT and JSPS (Japan), CNPq and FAPESP (Brazil), NSFC (China), MSMT (Czech Republic), IN2P3/CNRS and CEA (France), BMBF, DAAD, and AvH (Germany), OTKA (Hungary), DAE and DST (India), ISF (Israel), NRF and WCU (Korea), MES, RAS, and FAAE (Russia), VR and KAW (Sweden), U.S. CRDF for the FSU, US-Hungary Fulbright, and US-Israel BSF.

---

[*] Deceased

[†] PHENIX Spokesperson: jacak@skipper.physics.sunysb.edu

[1] M. J. Leitch et al. (FNAL E866/NuSea Collaboration), Phys. Rev. Lett. **84**, 3256 (2000).
[2] C. Lourenco, R. Vogt, and H. K. Woehri, JHEP **02**, 014 (2009), and references therein.
[3] D. de Florian and R. Sassot, Phys. Rev. **D69**, 074028 (2004).
[4] K. J. Eskola, V. J. Kolhinen, and R. Vogt, Nucl. Phys. **A696**, 729 (2001).
[5] D. Kharzeev and K. Tuchin, Nucl. Phys. **A770**, 40 (2006).
[6] A. Adare et al. (PHENIX Collaboration), Phys. Rev. Lett. **104**, 132301 (2010).
[7] T. Matsui and H. Satz, Phys. Lett. **B178**, 416 (1986).
[8] F. Karsch, D. Kharzeev, and H. Satz, Phys. Lett. **B637**, 75 (2006).
[9] A. Adare et al. (PHENIX Collaboration), Phys. Rev. **C77**, 024912 (2008).
[10] K. Adcox et al. (PHENIX Collaboration), Nucl. Instrum. Meth. **A499**, 469 (2003).
[11] A. Adare et al. (PHENIX Collaboration) to be published.
[12] S. N. White, AIP Conf. Proc. **792**, 527 (2005).
[13] A. Adare et al. (PHENIX Collaboration), Phys. Rev. **D82**, 012001 (2010).
[14] K. J. Eskola, H. Paukkunen, and C. A. Salgado, JHEP **04**, 065 (2009).
[15] R. Vogt, Phys. Rev. **C71**, 054902 (2005).
[16] D. Kharzeev and K. Tuchin, Nucl. Phys. **A735**, 248 (2004).
[17] S. R. Klein and R. Vogt, Phys. Rev. Lett. **91**, 142301 (2003).
[18] L. Frankfurt, V. Guzey, and M. Strikman, Phys. Rev. **D71**, 054001 (2005).
[19] J.-w. Qiu, J. P. Vary, and X.-f. Zhang, Phys. Rev. Lett. **88**, 232301 (2002).
[20] B. Z. Kopeliovich, I. K. Potashnikova, H. J. Pirner, and I. Schmidt (2010).
[21] M. B. Johnson et al., Phys. Rev. **C65**, 025203 (2002).
[22] L. A. Linden Levy, Nucl. Phys. **A830**, 353c (2009).